


LEGEND

RECENT	
LACUSTRIINE DEPOSITS	
24	Organic: muskeg, swamp, sedge bog, deep and wet organic deposits in depressional areas
23	Silt and clay: mixture of silt and clay, sand along some beaches, some organic materials
EROSIONAL DEPOSITS	
22	Slump: mixed glacial and bedrock materials; unstable slope
21	Gully, creek valley: thin colluvial cover over bedrock on valley slopes; thin alluvium materials along streams
ALLUVIAL DEPOSITS	
20	Alluvial fan: bedded silt, sand, and clay; variable thickness, overlying glacial deposits
19	Stream alluvium: mainly sand along major rivers; sand, silt, and clay along small streams
AEOLIAN DEPOSITS	
18	Aeolian sand: fine- to medium-grained sand in dunes and sheet form; developed from glacial outwash and deltaic deposits; thickest in dunes; thin to nearly absent between dunes
17	Loess: silt to very fine sand; thin to very thin
PLEISTOCENE	
GLACIOLACUSTRIINE DEPOSITS	
16	Clay and silt bedded clay and silt with minor sand; generally thick; level topography
15	Clay and silt containing pebbles and till-like layers; generally thin to very thin
14	Silt: massive to bedded silt with occasional pebbles; over five feet in thickness; level topography
13	Sand and silt: bedded sand and silt; generally thick; level topography with occasional small sand dunes
GLACIOFLUVIAL DEPOSITS	
12	Meltwater channel outwash: outwash sand and gravel occurring along the bottoms and terraces of meltwater channels; generally thin; level surface topography
11	Outwash sand: medium- to coarse-grained sand with pebbles and small gravel lenses; variable in thickness; gently undulating topography
10	Outwash sand and gravel: medium- to coarse-grained sand and gravel with some large boulders; variable in thickness; topography undulating
9	Kame, kame moraine: sand and gravel to gravel; variable in thickness but generally thick; rolling topography
GLACIAL DEPOSITS	
8	Colluviated ground moraine: till composed of sand, silt and clay with pebbles and boulders mantling colluviated steep slopes; partly bedded near surface; stable slope; generally thin
7	Gipsy till-hummocky moraine: predominantly sand derived from the Athabasca Sandstone; generally over forty feet thick; rolling topography
6	Gipsy till-ground moraine: predominantly sand derived from the Athabasca Sandstone; generally thin; undulating topography
5	Kinosis till-hummocky moraine: loam composition with numerous pebbles and boulders; generally thin; rolling topography
4	Kinosis till-ground moraine: loam composition with numerous pebbles and boulders; generally thin; undulating topography
3	Horse River till-hummocky moraine: silty clay composition; thick; rolling topography
2	Horse River till-ground moraine: silty clay composition; generally thin; undulating topography
DEVONIAN	
1	Methy Formation: dolomite forming numerous outcrops
Geological boundary: defined, approximate, assumed	
Abandoned beach	
Meltwater channel	
Meltwater channel scarp	
Esker	
Glacial flutings and drumlins	
Thin lacustrine deposit	


Geology by L.A. Bayrock and T.H.F. Reimchen, 1973

BEDROCK GEOLOGY



CRETACEOUS	
Kib	Labcie Formation: shale and silty shale; ironstone partings; silty fish-scale bearing beds
Kj	Joli Fou Formation: dark grey shale; silty interbeds
Kg	Grand Rapids Formation: quartzose and feldspathic sandstone; siltstone, silty shale; thin coal beds
Kc	Clearwater Formation: silty shale, siltstone, sandstone
Km	McMurray Formation: sandstone, siltstone, silty shale
DEVONIAN	
Dw	Waterways Formation: shale; argillaceous limestone
Dm	Middle Devonian: dolomite, limestone

Geological boundary
Surface contour (interval 500 feet)



SURFICIAL GEOLOGY
WATERWAYS
NTS 74D

LEGEND	
River or stream	
Intermittent river or stream	
Lake	
Intermittent lake	
Road, hard surface, all weather	
Railway	
Township boundary	
Section line	



DESCRIPTIVE NOTES

BEDROCK GEOLOGY

The Devonian strata of limestone, dolomite, shale, and evaporites underlie the entire area. The dolomites and dolomitic limestones of the Middle Devonian Methy Formation and the interbedded shale and limestone of the Upper Devonian Waterways Formation outcrop along the Athabasca and Clearwater Rivers, and their equivalents occur in the Bitumount area.

With the exception of the Labcie Formation, which is Upper and Lower Cretaceous, all other exposed bedrock formations are Lower Cretaceous. The McMurray Formation, which consists of sandstone, siltstone, and shale, lies unconformably on the Devonian limestones of the Waterways Formation and is well developed in the western part of the map area. The sandstone and shale are saturated with oil.

Tongues of shale in the upper part of the McMurray Formation are conformable with the overlying shales and sandstones of the Clearwater Formation.

The Grand Rapids Formation outcrops in the valleys and forms the first bedrock underlying the drift over much of the area.

The youngest rocks in the area, the Joli Fou and Labcie Formations, are marine shale.

SURFICIAL DEPOSITS AND LANDFORMS

Glacial Deposits

The area is one of extensive glaciation; till plains are divided into ground moraine and hummocky moraine plains, with the former existing as shallow, nearly level plains mantling the bedrock, and the latter as till plains with considerable local relief. In thickness, ground moraine is from 5 to close to 40 feet, and hummocky moraine is from 40 up to about 100 feet.

According to matrix composition, the tills of the area are divided into three main types: Gipsy till, which contains proportions of sand derived from the Athabasca Formation; the sand content exceeds 80 percent, and the till is made up by silt, pebbles, and boulders. In many locations the Gipsy till surface has been reworked by wind, which has produced small sand dunes. On cursory examination Gipsy till could be called outwash sand, and the landforms which have developed in it kames, but the silt content substantiates the fact that it is till.

Kinosi till is rich in gravel, having a matrix of a loam composition; that is, approximately equal proportions of sand, silt, and clay. It is present mainly in the Stony Mountain upland and north of the Clearwater River. On the Stony Mountain the till also contains a significant proportion of rounded gravel derived from Tertiary gravel, signifying that Tertiary gravel may still underlie the upland.

River till is a generally thin, clayey till containing fewer stones than the above two tills. It is present on the western flanks of Stony Mountain and north of the Athabasca River and underlies glaciolacustrine deposits over most of the area.

Glaciofluvial Deposits

Only one small esker is present in the area, located in the southeastern portion of the map area (Tp. 81, R. 2). The esker forms a sinuous ridge 20 feet high and 4 miles long, and consists of sand and gravel.

Kames are cone-shaped or rounded hills made of poorly sorted outwash sand and gravel; they may serve as a source of aggregate for construction purposes. Two or more closely associated kames are called a kame moraine. Generally the kames are steep-sided and 50 feet or more in height.

Sand and gravel outwash plains are relatively rare in the area. The largest one is located in township 83, range 11. Sand outwash plains consist of medium- to coarse-grained sand. The sand averages about 10 feet in thickness. Many of the sand outwash plains are covered with a variable thickness of sand and sand dunes and consequently are mapped as aeolian deposit areas.

The largest and most important potential gravel supply in the area comes from meltwater channel outwash, which is variable in thickness. Usually channel outwash is covered with a thin overburden of sand or even silt, so that testing for aggregate should extend to at least 10 feet, in order to penetrate the overburden.

Glaciolacustrine Deposits

Glaciolacustrine deposits cover a very large portion of the map area. On the basis of grain size composition, they have been divided into three groups of deposit sand, silt, and clay and silt.

Glaciolacustrine sand and silt deposits are generally present in the eastern half of the area. They are thin, ranging from a few feet to over 10 feet. The thin deposits are outlined on the map by a brown line symbol. They are generally less than 5 feet thick and overlie till.

Glaciolacustrine silt deposits are present only in the northwestern portion of the area. Their thickness ranges from a few feet to over 20 feet. Occasional pebbles occur in this deposit. The silt is very fine and is often very unstable; it is therefore prone to flowage, frost heaving, and slumping.

Glaciolacustrine clay and silt form the largest unit of lacustrine deposits in the area. The grain size ranges from loam to silty clay. Pure clay is very rare. The silty clay deposit (less than 5 feet) is also shown by a brown line symbol. In places, an irregular pattern in the northwest corner of the area, the clay and silt deposits alternate with significant proportions of pebbles and till-like lenses. These were formed primarily by drifting icebergs in glacial lakes. Generally, the pebbly pebbles may be so numerous that they give the lacustrine deposits a till-like appearance.

Aeolian Deposits

Loess, which is silt deposited by wind, is found only on the northern shore of Gordon Lake. It is massive and varies from silt to very fine sand in texture.

Aeolian sand occurs as dunes or as sheet sand. The dunes show effective wind direction from the southeast to east. Generally, the sand is medium-grained, massive, and very clean. In many locations where surface vegetation on the dunes has been stripped, boulders are present. The dunes vary from 10 to over 30 feet in height. Between the dunes, aeolian sand sheet deposits of variable thickness are present; generally they are very thin, ranging from about 2 to 5 feet thick.

Alluvial Deposits

Streams are still incising into bedrock, and streams have alluvium made of sand, silt, and clay. There is often a lag gravel layer deposited from the underlying till. The large rivers, on the other hand, have alluvium made of sand. Gravel-sized material is rare, and if present, it is of very local origin.

Alluvial fans are present mainly at the foot of Stony Mountain, Muskeg Mountain, and in the Clearwater River valley along the base of some steep banks. The material making up the alluvial fans is of sand to silty clay texture, generally thin, and bedded on a small scale.

Erosional Features

Gullies and creek valleys in the area have very steep slopes, and most of the glacial deposits are relatively thin. Thin colluvium mantles the slope, obscuring bedrock outcrops in most areas except where active erosion is taking place or where bedrock ledges protrude to the surface.

Stumps are the remains of mature trees on steep slopes. Generally the stump areas consist of a mixture of glacial and bedrock materials. The areas outlined as stump on the map are extremely unstable, and new movements of materials downslope can be expected.

Recent Lacustrine Deposits

Recent lacustrine deposits exposed above lake levels are rare, located around the edges in the eastern portion of the map area. Generally, they consist of silt and clay with organic material. Where former beaches were present, sand is encountered.

Organic deposits are present as swamps, muskegs, and peat bog and cover a very large portion of the map area. Generally they are very thin, ranging from 1 to 3 feet in depth. Deep bogs are relatively rare and are of the wet-sedge bog type. Only the deep sedge bogs have been delineated.

EXPLANATORY NOTE

Map users may note nomenclatural discrepancies between rock units on the Bitumount and Waterways map sheets. The discrepancies are explainable as follows. Gipsy till is composed essentially of sand with pebbles and boulders derived from the Athabasca Formation. In certain locations, specifically in the north-central corner of the Waterways area (NTS 74D), the Gipsy till is so devoid of sand that it has been projected across the area and supports occasional sand dunes. In the Bitumount area (NTS 74E) Gipsy till is well represented as outwash sand and gravel (legend unit 7), and as outwash sand and gravel mantled by glacier (legend unit 6). As new information became available, it was realized that these units actually are a till (Gipsy till) which in grain size composition approximates outwash.