

This is a common map legend for the surficial geology of northern Alberta. Coloured legend blocks indicate map units that appear on this map. Not all map symbols shown in the legend necessarily appear on this map.

UNIT SYMBOL	UNIT NAME	DESCRIPTION AND GENESIS
QUATERNARY		
HOLOCENE		
A	ANTHROPOGENIC MATERIALS:	Culturally-made or modified geological materials such that their physical properties (e.g., structure, cohesion, compaction) have been drastically altered.
O	ORGANIC DEPOSITS:	Undifferentiated black peat layers, woody to fibrous muck, occurring in undifferentiated wetlands, commonly underlain by fine-grained, poorly-sorted glaciolacustrine deposits, includes marshes, swamps, bogs and fens.
OB	ORGANIC BEDS:	Occurs in a peatland with a fluctuating water table and commonly a raised surface; peatland surface is dominated by sphagnum mosses, heath shrubs and short, stunted trees.
OF	Fan peat:	Occurs in a peatland with water table at surface and slow internal drainage; peatland surface is dominated by sedges, with grasses and reeds near local pools, and sparsely treed.
C	COLLUVIAL DEPOSITS:	Materials that have reached their present position as a result of direct, gravity-induced movement (commonly occurs as slope and slump deposits confined to valley slopes and floors); includes pre-existing bedrock, till, glaciolacustrine, glacioluvial and eolian sediments, generally poorly sorted.
F	FLUVIAL DEPOSITS:	Sediments transported and deposited by streams and rivers; synonymous with alluvial; includes well-sorted stratified sand, gravel, silt, clay and organic sediments occurring in channel and overbank deposits (e.g., oxbow, point-bar, terrace, fans and deltas).
L	LACUSTRINE DEPOSITS:	Sediments deposited in and adjacent to recent lakes; offshore sand, silt and clay, minor organic deposits; littoral (nearshore beaches and bars) sand and silt and minor gravel.
E	EOLIAN DEPOSITS:	Wind-deposited sediments, well-sorted, medium- to fine-grained sand, and minor silt (loess); generally massive to locally cross-bedded or ripple laminated; includes both active and vegetated deposits.
PLEISTOCENE		
LG	GLACIOLACUSTRINE DEPOSITS:	Fine-grained distal sediments deposited in or along the margins of glacial lakes, including varved clays that were reworked by the melting of floating ice; includes laminated (rythmically bedded) to massive fine sand, silt and clay, and may contain ice-melted stones.
LGL	Lateral and nearshore sediments:	Massive to stratified well-sorted silt, sand and minor gravel; occurs as beaches, bars, spits and forest detrital deposits deposited during regression and lowering of glacial lakes.
FG	GLACIOLUVIAL DEPOSITS:	Sediments deposited by glacial meltwater streams directly in front of glacier ice as subglacial or subaqueous outwash; includes sand and gravel, often stratified, minor silt, and may show evidence of ice melting (slumped structures). Features include meltwater channels, kettle holes and terraces.
FOG	Ice-contact sediments:	Sediments deposited by glacial meltwater streams in direct contact with glacial ice, either in front of (fame terraces) or within glacial ice (eskers, crevasse ridges); includes massive to stratified, poor to moderately sorted coarse sediments (predominantly pebble gravel and coarse sand, locally till) and may show evidence of ice melting (slumped structures).
M	MORANE:	Material deposited directly by glacial ice (without modification by any other agent of transportation); includes nonsorted glacial till as lodgement till (a mixture of silt, clay, sand and minor pebbles, cobbles and boulders) at the ice margin or beneath a glacier. Locally, it may contain blocks of bedrock, pre-existing stratified drift and till. Beds and lenses of glaciolacustrine and glacioluvial sediments may occur.
MS	Stagnant ice moraine:	Terrain resulting from the collapse and lateral movement of englacial and supraglacial sediment in response to melting of buried stagnant ice at the ice margin; sediment is mainly detrital (till), but locally includes stratified sediments of glaciolacustrine or glacioluvial origin. Characterized by low- to high-relief hummocky topography.
MT	Ice-thrust moraine:	Terrain resulting from glacio-tectonic transport of originally subglacial sediment and by the glacier more or less intact; deposits may include syngenetic till as well as masses of deposited pre-existing till, stratified drift and/or bedrock. Characterized by high to moderate relief and features include hill-hole pairs and glacio-tectonic moraine ridges.
MF	Fluted moraine:	Glacially streamlined terrain; varies from alternating furrows and ridges to nearly equidimensional smoothed hills; all landforms parallel to the local ice flow direction; includes flutes, chertins and drumlins.
ML	Undifferentiated moraine:	Moraine of undetermined origin and texture, but usually fine-grained; silt, clay and minor sand; lacks distinctive features required to subdivide it to the unit as a composite containing three or more subdivisions inseparable at the map scale.
PRE-QUATERNARY		
R	BEDROCK:	Undivided; may include crystalline (Shield), carbonate or clastic sedimentary rock, and/or coal.
RT	Tertiary gravels:	Predominantly quartzite and chert gravel and cobbles; preglacial age.

FEATURES LEGEND


Thermokarst depression	⊖
Landslide and active layer failure scar (small)	⊖
Landslide and active layer failure scar (large)	⊖
Eolian forms, dune ridges	⊖
Beach or strandline	⊖
Wave cut bench	⊖
Escarpment	⊖
Meltwater channel (minor)	⊖
Meltwater channel (minor, flow indicated)	⊖
Meltwater channel (major)	⊖
Meltwater channel (major, flow indicated)	⊖
Crevasse filling	⊖
Ice contact slope	⊖
Kettle	⊖
Esker, direction of paleoflow unknown	⊖
Esker, direction of paleoflow indicated	⊖
Drumlinoid or streamlined landform	⊖
Drumlinoid, down-ice flow indicated	⊖
Buried drumlinoid or streamlined landform	⊖
Minor moraine ridge: De Geer, Rogen, ribbed, washboard (minor)	⊖
Major moraine ridge	⊖
Ice thrust ridge	⊖
Stratification (direction unknown)	⊖
Stratification (direction known)	⊖
Bedrock outcrop	⊖
Gravel and/or sand pit	⊖

ROADS LEGEND

Paved	⊖
Gravel	⊖
Unimproved	⊖
Trail	⊖
UTM, Zone 11 Grid	⊖
Contour, intervals 10 metres	⊖

UNIT NOTATION

Example: GLACIOLACUSTRINE plain



Textural Modifier
Textural characteristics may be applied to the terrain classification as a prefix based on field observations or by inference from distinctive genesis and/or morphology. When two modifiers are given, the second letter is the dominant texture, with the first letter indicating the secondary texture, i.e., sc for sandy clay.

GENETIC & GEOMORPHIC MODIFIERS

c	crevasse fill	Ice-contact ridges and linear forms deposited by meltwater in stagnant ice
d	doughnut ridges and ridges	arcuate hummocks with a central depression (doughnut ridges), plateau mounds and brain pattern ridges, low to moderate relief
e	eroded	planar surface eroded by glacial meltwater, often capped by a boulder lag deposit and/or thin deposit of sand and gravel
f	fan	gently sloping fan-shaped mass of detrital debris
g	gullied	slopes dissected by modern ravines created by intermittent runoff
h	hummock	assemblage of approximately equidimensional hills and hollows; moderate to high relief (commonly greater than 2 m)
k	collapse	depression, including kettles, tilted outwash, thermokarst depressions, karst sinkholes
m	meander	sinuous curves, loops and oxbows produced as meltwater and modern streams shift their channel over time
p	plain	deposit greater than 2 m thick; commonly masks geomorphic pattern of underlying deposits; flat to gently rolling topography (commonly less than 2 m relief)
r	ridged	one or more parallel or subparallel, convex, linear morphological elements with a width-to-length ratio greater than 2; low to high relief
s	slumped	landslide blocks, slope failure debris
t	terrace	terrace bench cut by either meltwater or wave action; antiplication terrace, kame terrace
u	undulating	low-relief rolling terrain, well and scale topography
v	veneer	thin mantle of unconsolidated material too thin to mask the minor irregularities of the surface of the underlying material. It ranges in thickness from 10 cm to 1 metre and may be discontinuous.
w	washboard	low relief transverse moraine ridges, usually formed from basal ice shearing
z	dissected	channelled or dissected by glacial meltwater flow, dissected terrain by Holocene fluvial activity
y	delta	lake delta; ice-contact delta

Complex

Where two or more classes of terrain are interspersed in a mosaic or repeating pattern on a scale too small to warrant meaningful differentiation, the proportion of each component in the complex is given in a two or three position designation set off by slashes denoting arbitrary percentage limits. For example,

MpL.Gv means that the area is underlain by approximately 60% moraine plain and up to 40% glaciolacustrine veneer.

MvL.GvF.Gv means that at least 60% of the area is underlain by moraine veneer, with up to 40% glaciolacustrine veneer and less than 10% glacioluvial plain.

L.Gv/M means that more than 60% of the area is underlain by a glaciolacustrine plain, with less than 15% moraine.

Stratigraphic Sequence

Where materials of different origin or texture are known to be superimposed or can be confidently inferred, the sequence is indicated in conventional order using vertical separators, such as:

L.Gv / M / This sandy glaciolacustrine sediment deposited on moraine plain


Transitional Association

Locally, two or more terrain units are juxtaposed by means of related origin, temporal sequence, or ambiguous geomorphic distinction. In the latter case, both components may or may not be present. Such situations are identified by a compound designation marked by a hyphen. Examples are: "FGz-LG" indicating ice-contact delta indistinguishable from glaciolacustrine delta, or "FG-Mv" indicating ice-contact kame and kettle topography that blends with hummocky stagnant ice moraine.


Morphologic Overprint

Where a sequence of geomorphic processes has produced a multi-aspect or compound terrain fabric, the geomorphic modifier suffixes are appended in the inferred order of superposition. "Mry" means that a plain of fill has been modified into ridge forms and finally dissected by modern streams. "FGgr" means that a glacioluvial plain has been discontinuously covered by ice-contact hummocks and ridges.

Bedrock Topography of NTS 84B (Peelers Lake)



Drift Thickness of NTS 84B (Peelers Lake)



Acknowledgements:

Initial fieldwork was carried out by the Alberta Geological Survey (AGS) during the summers of 1997, 1998 and 1999. Surficial mapping was completed in 2001 under the AGS's Quaternary mapping initiative. Jill Weiss and Charlene Brighent provided assistance with fieldwork and compilation of the digital databases. Digital cartography and GIS was completed by Monica Price, Joan Waters, Jill Weiss and Natasha Blundon.

Digital base produced by the Resource Data Division, Alberta Environment, supplied by Spatial Data Warehouse Ltd.

References:

Croft, W. J. (1979). Hydrogeology of the Peelers Lake area, Alberta. Alberta Energy and Utilities Board, EUB/AGS Earth Sciences Report 79-05, 120.

Lindsay, J. D., Hering, F. K., Pawlik, S. and Ostry, W. (1957). Exploratory soil survey of Alberta map sheets 84C (west half), 84B (84A and 74D), Alberta Exploratory Soil Survey Report No. 58-1, 36 pp.

Pawlowicz, J. G. and Fenton, M.M. (2002). Bedrock topography of the Peelers Lake map area (NTS 84B). Alberta Energy and Utilities Board, EUB/AGS Map 252.

Pawlowicz, J. G. and Fenton, M.M. (2003). Drift thickness of the Peelers Lake map area (NTS 84B). Alberta Energy and Utilities Board, EUB/AGS Map 253.

Pawlowicz, J. G., Dufresne, M.B. and Fenton, M.M. (1998). Diamond indicator minerals from auger core holes, a possible second dispersal train in the Peelers Lake map area (84B). Alberta Energy and Utilities Board, EUB/AGS Geoscience 98-02.

Pawlowicz, J. G. and Fenton, M.M. (1998). Auger core lithology from the Wabasca/Red Earth area, north-western Alberta. Alberta Energy and Utilities Board, EUB/AGS Geoscience 98-02.

Pawlowicz, J. G., Meeks, J.O. and Fenton, M.M. (1999). Auger core descriptions, 1998 drilling, Peelers Lake map sheet (NTS 84B), North-central Alberta, Alberta Energy and Utilities Board, EUB/AGS Open File Report 99-02.

Scarlett, W., Green, P.C., Blundon, W. and Barthelemy, H. (1988). Aggregate resources Peelers Lake 84B, Alberta Geological Survey Map A98B.

Published 2003
Copies of this map may be obtained from:
Information Sales
Alberta Geological Survey
Telephone: (781) 422-3767
Web site: www.agp.gov.ab.ca

Map 266
Surficial Geology of the Lubicon Lake Area, Alberta (NTS 84B/SW)

Geology by: M.M. Fenton, R.C. Paulen and J.G. Pawlowicz

Scale 1:100 000

Projection: Universal Transverse Mercator, Central Meridian 115° 30'
Datum: North American Datum, 1983

