



This legend is common to maps OF5460 and OF5461. Coloured legend blocks indicate map units that appear on this map. Not all map symbols shown in the legend appear on this map.

Note: In areas where the surficial cover forms a complex pattern, the area is coloured according to the dominant unit and labelled in descending order of cover (e.g. Q-Ts). Where buried aggregate deposits (sand and gravel - commonly associated with Gl surficial units) are known, or suspected, areas are coloured according to the overlying unit and labelled in the following manner: Lx/Gl.

QUATERNARY SURFICIAL DEPOSITS

- POST-LAST GLACIATION**
- NONGLACIAL ENVIRONMENTS**
- ANTHROPOGENIC DEPOSITS:** culturally-made or modified geological materials such that their original physical properties (e.g. structure, cohesion, compaction) have been drastically altered; >2 m thick.
- ORGANIC DEPOSITS:** peat and muck, 1 to 3 m thick on average, formed by the accumulation of plant material in various stages of decomposition; generally occurs as flat, wet terrain (swamps and bogs) over poorly drained substrates.
- Bog peat:** sphagnum or forest peat formed in an ombrotrophic environment; wet terrain; may be treed or treeless; O'h, hummocky, mounds and plateaus; area may be undrained by ground ice or shallow permafrost conditions.
- Fen peat:** peat derived from sedges and partially decayed shrubs in a eutrophic environment; forms relatively open peatlands with a mineral-rich water table that persists seasonally near the surface; generally covered with low shrubs and an occasional sparse cover of trees.
- Undifferentiated hummocky bog and fen deposits.**

- COLLUVIAL DEPOSITS:** mass wasting debris; poorly sorted, massive to stratified debris deposited by direct, gravity-induced movement; composition dependent on source.
- Landslide and slump debris:** active and inactive landslides; hummocky topography; clast size: generally >1 to 10 m thick, but may exceed 10 m near the toe of large landslides.
- Colluvial veneer:** thin and discontinuous cover of slumped and/or soliflucted material <1 m thick, overlies bedrock or till.
- ALLUVIAL DEPOSITS:** sorted gravel, sand, minor silt, and organic debris deposited by streams; commonly stratified.
- Floodplain deposits:** sorted gravel, sand, silt, and organic debris >1 m thick; forming active floodplains close to river level with meander channels and scroll marks.
- Fluvial terrace deposits:** inactive terraces above modern floodplain; >2 m thick; represents a potential aggregate source.
- Alluvial fan deposits:** poorly sorted gravel, sand, and organic debris >1 m thick.
- Undifferentiated fluvial deposits.**


- LACUSTRINE DEPOSITS:** sand, silt, and minor clay deposited in a former lake; >1 m thick; generally overlain by organic deposits; exposed by recent fluctuations in lake levels.
- NONGLACIAL AND PROGLACIAL ENVIRONMENTS**
- EOLIAN DEPOSITS:** wind-deposited medium to fine sand; derived from deltaic or glaciolacustrine deposits; in some areas eolian sediments are thin or absent between dunes.
- Ridged eolian deposits:** forming dunes; generally >2 m thick.

- POSTGLACIAL OR LATE WISCONSINAN**
- PROGLACIAL AND GLACIAL ENVIRONMENTS**
- GLACIOLACUSTRINE DEPOSITS:** fine sand, silt, and clay, with minor debris-flow diamict, deposited in glacially-dammed lakes in valleys and along the margin of the retreating Laurentide Ice Sheet; usually overlain by organic deposits in lowlands.
- Glaciolacustrine blanket:** >1 m thick; Lbh, hummocky glaciolacustrine sediments; >1 m thick; forming circular hummocks and hills surrounded by depressions with a relief usually >2 m; interspersed with minor fans.
- Glaciolacustrine veneer:** thin and discontinuous; <1 m thick.

- GLACIOFLUVIAL DEPOSITS:** well to poorly stratified sand and gravel; minor diamict; deposited behind, at, or in front of the ice margin by glacial meltwater; represents a potential aggregate source.
- Proglacial outwash:** cross-stratified gravel and sand deposited in front of the ice margin; Gl; outwash plain deposits; generally 1 to 5 m thick; generally mantle valley floors and surfaces adjacent to glacial meltwater channel margins; Gl; outwash terrace deposits, generally associated with meltwater channels and canyons; 1 to 10 m thick.
- Ice-contact stratified drift:** poorly sorted sand and gravel with minor diamicts; deposited in contact with the retreating glacier; 1 to >20 m thick; Gh, hummocky topography relating to melting of underlying ice; Gl; esker ridges.

- TILL:** diamict deposited directly by the Laurentide Ice Sheet; sandy to clayey matrix with stratified clasts of various lithologies, including many Canadian Shield, carbonate, and sandstone erratics; clast content is typically low (<10 %).
- Till blanket:** >1 m thick, continuous till cover forming undulating topography that locally obscures underlying units.
- Streamlined and fluted till:** >1 m thick, till surface marked by streamlined landforms including flutes and drumlins.
- Hummocky till:** >1 m thick; hummocky till surface.
- Ridged till deposits:** >1 m thick, moraines or crevasse fillings forming a ridged topography.
- Till veneer:** <1 m thick, discontinuous till cover; underlying bedrock topography is discernible.

- PRE-QUATERNARY BEDROCK**
- Sedimentary bedrock:** Cretaceous Fort St. John Group shales (including the Shalshbury Formation) and Dungeness Formation sandstone exposed in highlands and along meltwater channel and canyon walls.



DESCRIPTIVE NOTES

The Caribou Creek map area is located in northwestern Alberta within the Fort Nelson and Peace River lowlands (Bostock, 1987). The Fort Nelson Lowlands is a region of flat relief with an elevation varying from 300 to 500 m above sea level and is incised by the Peace River. The Peace River lowlands, through the Peace River drainage, to the east of the Peace River which is part of the Mackenzie River drainage. The main economic activities in the map area include agriculture and logging and to a lesser extent oil and gas.

The surficial geology of the map area was interpreted from 1:50 000 scale black and white air photographs dating from 1960 and produced by the Alberta Sustainable Resource Development. To establish the surficial geology, the surficial geology was interpreted from the Shuttle Radar Topography Mission (SRTM) imagery (SRTM 3-arc second, 30 m resolution). Field work was conducted during a 3 week period in the summer of 2006. The collection of field observations necessary for the mapping of the surficial geology, the logging of stratigraphy exposed in pits and river bank exposures, and the collection of bulk glacial landform samples (approx. 50 kg per sample). Access to the region was by truck, all terrain vehicle, and foot travel. A hand auger was utilized to identify the sediment within the surface due to the paucity of natural and man-made sediment exposures. The surficial geology was interpreted on air photographs and was subsequently digitized by a consultant using a digital vector plot (DVP) system.

The bedrock of the map area consists of horizontally bedded, poorly indurated shale of the Shalshbury Formation (part of the Fort St. John River Formation) (Green et al., 1970; Chalko, 2008). The only bedrock exposure in the map area is located along the north shore of the Peace River and consist of horizontally bedded, poorly indurated shale of the Shalshbury Formation (Chalko, 2008).

Till in the Caribou Creek map area consists of a clayey diamict (10 to 40% clay) with a clast content varying from 10 to 20 %. Clast lithologies in till consist of locally derived shale and limestone interbedded with Canadian Shield and Proterozoic Paleozoic sedimentary rocks transported from the northwest.

Glaciolacustrine sediments are rare. One esker was mapped in the south east part of the map area, from an air photo interpretation with no ground evaluation. The esker formed during glacial conditions and subsequently was submerged in a glacial lake early during deglaciation. This esker was mapped as an ice-contact glaciolacustrine deposit, but may be overlain by a veneer of glacial lake sediments.

Glacial lake sediments are dominantly massive and rarely bedded. Because of the low clast content of till and poor sediment exposures the distinction between till and glacial lake sediments was difficult in several places. Glacial lake sediments generally thin to the west with respect to present surface elevation. The maximum elevation reached by glacial lake sediments is estimated to be 410 m. Glacial lake sediment cover is discontinuous near the contact zone with till (in present and in the southwest sector of the map area). Therefore, the glacial lake which formed in the region reached a minimum elevation of 410 m. Glacial lake sediment cover is discontinuous near the contact zone with till. Certain regions below the maximum extent of glacial lake sediment have been mapped as till, as opposed to glacial lake sediments, because of the high density and good definition of the glacial flutes. In certain areas hundreds of meters in diameter with a relief >2 m in relief (Lbh). These landforms are interpreted to result from periglacial processes whereby local upthrust occurred because of freezing of thawed layers immediately following rapid glacial lake drainage. Similar features were observed in the surrounding regions to the south, within the Sturgeon Lake map area (Henderson, 1999).

Organic deposits in the form of fens and bogs are omnipresent in the Caribou Creek map area. Permafrost is still present in the hummocky bogs (Lbh).

Fluting and drumlins ridges are well developed in the region. Within the Caribou Creek map area, fluting morphologies are apparent through the cover of glacial lake sediments and organic deposits. In other words, the fluting are draped by a veneer of glacial lake sediments or organic deposits but are still visible on air photographs. Fluting are a key indicator of ice-flow directions in this region where bedrock striations are absent because of the poorly indurated nature of the bedrock. Only minor moraines have been mapped in the region. Moraines, undeniably, retreat to the east-northeast. Circular landforms of unknown origin were observed in the central and east central part of the map area. Some of these features are composed of multiple, tightly spaced and very low relief circular ridges. Their origin glacial, or periglacial is still unclear. Similar features are present to the north (Plouffe et al., in press).

The Laurentide Ice Sheet, or periglacial is still unclear. Similar features are present to the north (Plouffe et al., in press). The Laurentide Ice Sheet advanced westward to the Caribou Creek map area. At glacial maximum, ice was generally found to the southwest as evidenced by the orientation of fluting and drumlin ridges. Retreat of ice from the area occurred between 11 500 and 11 000 radiocarbon years BP (13 450 to 13 000 calendar years BP) (Dyke, 2004). The southwestern sector of the map area was deglaciated by the Laurentide Ice Sheet. The Laurentide Ice Sheet was rooted in the underlying sediments and served as the outlet of the Keg River stage of glacial lake Peace (Matthews, 1980). Glacial lake Peace expanded to the east, ice retreated easterly and responded the eastward drainage. The extent and elevation of this glacial lake diminished as ice retreated easterly. At some point, the level of glacial lake Peace was controlled by an outlet meltwater flow spilling at an elevation of 300 m and immediately north of the map area (Plouffe et al., in press). The Peace River drainage within the Caribou Creek map area was established following the eastward migration of glacial lake Peace above the retreating ice front. The river went through a cycle of aggradation and degradation leaving thick accumulation of sand and gravel in the form of buried terraces. During early post-glacial time, the development of an eastern vegetation cover, westerly winds reworked part of the sandy sediments into eolian dunes. Following deglaciation, vegetation invaded the region and eolian dunes were accumulated. Continued eolian accumulation of sand and gravel in the hummocky terraces allowed for the development of permafrost in northwestern Alberta (Zolal, 1993). Permafrost is still present in the hummocky terraces (Lbh). Because of the generally flat topography of the region, the continuous damming of small streams by layers, the energy of secondary streams is low and time fluvial incision has occurred.

Fluvial terraces are limited in this region. The Laurentide Ice Sheet has always been present in the fluvial terraces of the Peace River. Other fluvial deposits in the area are unlikely to represent potential aggregate sources. For information on aggregate sources, see the Caribou Creek map area (Plouffe et al., 2007a, 2007b, and to the east (Plouffe and Plouffe, 2007b). Capable bed assistance was provided by Thomas Anwarachuk, Ervin Fournier, Chris Kowalski, Thomas Talley, and Kelly Lee Wang Goss.

REFERENCES

Bostock, H.S.
1987. Physiographic regions of Canada; Geology, Map 1254A, scale 1:5 000 000.

David, P.R.
1972. Sand dune occurrences of Canada: a theme and resource inventory study of eolian landforms of Canada. Indian and Northern Affairs, National Parks Branch, Ottawa, Contract 14-200, 180 p.

Dyke, A.S.
2004. An outline of North American deglaciation with emphasis on central and northern Canada; in Quaternary glaciations - extent and chronology, part II, (eds.) J. Ehlers and P.L. Gibbard, Elsevier, Amsterdam, p. 373-424.

Edwards, D., Budney, H., Berezniak, T., and Burkolc, L.
2004. Sand and gravel deposits with aggregate potential, Mount Waut, Alberta (NTS 84K), Alberta Energy and Utilities Board, EUB/AGS, Map 308, scale 1:250 000.

Green, R., Melton, G.B., and Carigny, M.A.
1970. Bedrock geology of northern Alberta, Research Council of Alberta, scale 1:500 000.

Henderson, E.R.
1998. Surficial geology of Sturgeon Lake map area, Alberta; Geological Survey of Canada, Ottawa, Memoir 303, 108 p.

Matthews, W.H.
1980. Retreat of the last ice sheets in northeastern British Columbia and adjacent Alberta; Geological Survey of Canada, Bulletin 321, 22 p.

Oulitch, A.
2008. Bedrock geology, Peace River, Alberta; Geological Survey of Canada, Open File 5285, scale 1:1 000 000.

Paulen, R.C., Fenon, M.M., Weiss, J.A., Pawlowicz, J.G., Plouffe, A., and Smith, I.R.
2005. Surficial geology of the Hay Lake area, Alberta (NTS 84LNE); Alberta Energy and Utilities Board, EUB/AGS Map 375, scale 1:100 000.

Paulen, R.C. and Plouffe, A.
2007a. Surficial geology of the Buabe River-Riverton River area (NTS 84KNE); Alberta Energy and Utilities Board, EUB/AGS Map 473 and Geological Survey of Canada, Open File 5255, scale 1:100 000.

Paulen, R.C. and Plouffe, A.
2007b. Surficial geology of the La Cible area (NTS 84KSE); Alberta Energy and Utilities Board, EUB/AGS Map 472 and Geological Survey of Canada, Open File 5256, scale 1:100 000.

Plouffe, Z.K. and Wolfe, S.A.
2002. Sand dune orientations in the prairie provinces of Canada; Geological Survey of Canada, Ottawa, Open File 417, CD-ROM.

Plouffe, A., Kowalski, C.J., and Paulen, R.C.
2007. Surficial geology, Meander River, Alberta (NTS 84KNV); Geological Survey of Canada, Open File 5461, Alberta Energy and Utilities Board, Alberta Geological Survey, File 414, scale 1:100 000.

Paulen, R.C., Fenon, M.M., Weiss, J.A., Pawlowicz, J.G., Plouffe, A., and Smith, I.R.
2004. Surficial geology, Bassett Lake, Alberta (NTS 84L SE); Geological Survey of Canada, Open File 4637, scale 1:100 000.

Zolal, S.C.
1993. Cyclic development of permafrost in the peatlands of northwestern Alberta, Canada, Arctic and Alpine Research, v. 25, p. 240-246.

GSC OPEN FILE 5460
EUB/AGS MAP 415
SURFICIAL GEOLOGY
CARIBOU CREEK
ALBERTA

Authors: A. Plouffe and R.C. Paulen
Geology by A. Plouffe, 2005-2006
Airphoto interpretation by A. Plouffe, 2005-2006
Digitizing and digital cartography by Geotech, Geomatics services
Digital map compilation by L. Robertson, Northern Canada Division, 2006
Digital cartography by J.D. Narraway, Data Dissemination Division (DDD)

Scale 1:100 000/Echelle 1/100 000
kilometres 2 4 6 8 kilometres

Universal Transverse Mercator Projection
North American Datum 1983
© Sa Majesté la Reine du chef du Canada 2007

Projection transversale universelle de Mercator
Système de référence géodésique nord-américain, 1983
© Sa Majesté la Reine du chef du Canada 2007

This map was produced from processes that conform to the Scientific and Technical Publishing Services Subdivision (DDD) Quality Management System, registered to the ISO 9001:2000 standard.

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada.

Digital base map from data compiled by Alberta Sustainable Resource Development, modified by DDD

Digital base map from data compiled by Alberta Sustainable Resource Development, modified by DDD

Mean magnetic declination 2007, 20'03"E, decreasing 2.7" annually. Readings vary from 20'25"E in the NW corner to 19'41"E in the SE corner of the map.

Elevations in metres above sea level

OF5670	B4 M	B4 N	B4 O
OF5237	OF5184	B4 L	B4 J
OF4754	OF4637	OF5460	OF5629
B4 E	B4 F	B4 G	B4 G

OPEN FILE DOSSIER PUBLIC
5460
GEOLOGICAL SURVEY OF CANADA / COMMISSION GÉOLOGIQUE DU CANADA
2007

Open files are products that have not gone through the GSC formal publication process.
Les dossiers publics sont des produits qui n'ont pas été soumis au processus de publication de la GSC.

Recommended citation:
Plouffe, A. and Paulen, R.C.
2007. Surficial geology, Caribou Creek, Alberta; Geological Survey of Canada, Open File 5460; Alberta Energy and Utilities Board, Alberta Geological Survey, Map 415, scale 1:100 000.

