


This is a common map legend for the surficial geology of northern Alberta. Coloured legend blocks indicate map units that appear on this map. Not all map symbols shown in the legend necessarily appear on this map.

UNIT	UNIT NAME	DESCRIPTION AND GENESIS
QUATERNARY		
Holocene		
A	Anthropogenic materials	Artificially made ground or geological materials that have been disturbed by human activity, such that their physical properties (e.g. structure, cohesion, compaction) have been drastically altered.
O	Organic deposits	Undifferentiated peat (woody to fibrous muck) occurring in wetlands; commonly underlain by fine-grained, poorly drained glaciolacustrine deposits; includes marshes, swamps, bogs and fens.
OB	Bog peat	Occurs in a peatland with a fluctuating water table and commonly a raised surface; peatland surface is dominated by sphagnum mosses, heath shrubs and short, stunted trees.
OF	Fen peat	Occurs in peatland which receives water from slowly flowing streams and groundwater, with the water table lying at the land surface; peatland surface is dominated by sedges, with grasses and reeds near local pools, and is sparsely treed.
C	Colluvial deposits	Materials that have reached their present position as a result of direct, gravity-induced movement; commonly occurs as slope and slump deposits confined to valley slopes and floors; includes pre-existing bedrock, till, glaciolacustrine, glaciofluvial and eolian sediments, generally poorly sorted.
F	Fluvial deposits	Sediments transported and deposited by streams and rivers; synonymous with alluvium. Includes well-sorted stratified sand, gravel, silt, clay and organic sediments occurring in channel and overbank deposits (e.g. postglacial floodplains, terraces, fans and deltas).
L	Lacustrine deposits	Sediments deposited in and adjacent to recent and modern lakes; includes offshore sand, silt and clay, minor organic deposits, may also include minor littoral (nearshore) beaches and bars composed of sand, silt and minor gravel.
E	Eolian deposits	Wind deposited sediments; comprise well-sorted, medium- to fine-grained sand and minor silt; generally massive to locally cross-bedded or ripple-laminated; includes both active and vegetated dunes and sand sheets.
Pleistocene		
LG	Glaciolacustrine deposits	Primarily fine-grained, distal sediments deposited in or along the margins of glacial lakes, including sediments released by the melting of floating ice. Includes laminated (hythmically bedded) to massive fine sand, silt and clay, and may contain ice-rafted debris.
LGL	Littoral and nearshore sediments	Massive to stratified, well-sorted silty sand, pebbly sand and minor gravel; occurs in beaches, bars, spits and deltas; deposited during regression and lowering of glacial lakes.
FG	Glaciofluvial deposits	Sediments deposited by glacial meltwater streams as subaerial or subaqueous outwash. Includes sand and gravel, often stratified, minor silt, and may show evidence of ice melting (slumped structures). Features include meltwater channels, kettle holes, terraces and minor ice-contact sediments.
FGI	Ice-contact sediments	Sediments deposited by meltwater streams flowing either in direct contact with the ice margin (kame terraces) or within and/or under glacial ice (eskers, crevasse ridges). Includes massive to stratified, poor to moderately sorted, coarse-grained sediments (predominantly pebble gravel and coarse-grained sand, locally till) and may show evidence of ice melting (slumped structures).
M	Moraine	Diamictic (ill) deposited directly by glacial ice and consisting of a mixture of clay, silt, sand and minor pebbles, cobbles and boulders. Locally, this unit may contain blocks of bedrock, pre-existing stratified sediment and till, or lenses of glaciolacustrine and/or glaciofluvial sediment.
MS	Stagnant ice moraine	Material resulting from the collapse and slumping of englacial and supraglacial sediment in response to the melting of buried stagnant ice at the ice margin; sediment is mainly diamictic, but locally includes stratified sediments of glaciolacustrine or glaciofluvial origin. Characterized by low to high-relief hummocky topography.
MT	Ice-thrust moraine	Terrain formed from the glaciostatic displacement of materials as blocks or rafts in a more or less intact state. Materials may include syngenetic till, as well as masses of pre-existing sediments and/or bedrock. Characterized by low to moderate relief and features include hill-hole pairs and glaciostatic moraines.
MF	Fluted moraine	Glacially streamlined terrain; varies from alternating furrows and ridges to nearly equidimensional smoothed hills; all landforms parallel the local ice flow direction; includes flutes, drumlins and drumlinoids.
FP	Pre-glacial fluvial deposits	Sediments transported and deposited by streams and rivers prior to glaciation. This includes sand and gravel deposited in paleovalleys (i.e., preglacial floodplains, terraces, fans and deltas).
PRE-QUATERNARY		
RT	Unconsolidated fluvial gravels	Predominantly well-sorted, quartzite and chert gravel and cobbles; Cordilleran source, Paleogene to Neogene.
R	Bedrock	


SYMBOL LEGEND		BASEMAP LEGEND	
Eolian forms, dune ridges		Primary road, paved	
Beach or strandline		Primary road, gravel	
Wave-cut bench		Unimproved road	
Escarpment		Truck trail	
Meltwater channel (minor)		River	
Crevasse filling		Lake	
Ice-thrust ridge		UTM, Zone 11 Grid	
Drumlinoid or streamlined landform		Contour, intervals 50 metres	
Buried drumlinoid or streamlined landform		Town	
Minor moraine ridge			
Major moraine ridge			
Iceberg scour			

UNIT NOTATION			
Example: sandy GLACIOLACUSTRINE plain			
	Textural modifier	Geomorphic unit	Unit
Textural Modifier			
Textural characteristics may be applied to the terrain classification as a prefix based on field observations or by inference from distinctive genesis and/or morphology. When two modifiers are given, the second letter is the dominant texture, with the first letter indicating the secondary texture, i.e., sc for sandy clay.			
p	pebble		
g	gravel		
s	sand		
\$	silt		
c	clay		
a	sand-silt-clay		
GENETIC & GEOMORPHOLOGICAL MODIFIERS			
c	crevasse fill	ice-contact ridges formed by the slumping of sediment into crevasses on the ice surface or the squeezing of till into fractures at the ice base	
d	doughnut rings and ridges	circular hummocks with a central depression, plateau mounds and brain-like pattern ridges, low to moderate relief	
e	eroded	planar surface eroded by glacial meltwater, often capped by a boulder lag and/or thin deposit of sand and gravel	
f	fan	gently sloping fan-shaped mass of detrital debris	
g	gullied	slopes dissected by modern ravines created by intermittent runoff	
h	hummock	assemblage of approximately equidimensional hills and hollows; moderate to high relief (commonly greater than 2 m)	
k	collapse	depression, includes kettle holes, pitted morphology, thermokarst depressions, karst sinkholes	
m	meander	sinuous curves, loops and oxbows produced as meltwater and modern streams shift their channels over time	
p	plain	deposit greater than 2 m thick, commonly masks geomorphic pattern of underlying deposits; flat to gently rolling topography (commonly less than 2 m relief)	
r	ridged	one or more parallel or subparallel, convex, linear morphological elements with a length-to-width ratio greater than 2; low to high relief	
s	slumped	landslide blocks, slope failure debris	
t	terrace	a bench of either erosional or depositional origin that flanks the sides of floodplains, valleys and lakes; includes fluvial and glaciofluvial terraces, shoreline terraces and angiplation terraces	
u	undulating	low-relief rolling terrain; swell and swale topography	
v	venerer	thin mantle of unconsolidated sediment that is too thin to mask the minor irregularities of the surface of the underlying material; it ranges in thickness from 10 cm to 1 metre and may be discontinuous	
y	dissected	channelled or dissected by glacial meltwater and/or Holocene fluvial activity	
z	delta	lake delta; ice-contact delta	
Complex			
Where two or more classes of terrain are interspersed in a mosaic or repeating pattern on a scale too small to warrant meaningful differentiation, the proportion of each component in the combination is given in a two- or three-position designation set off by slashes denoting arbitrary percentage limits. Examples are:			
'MpLgV'	indicates the area is underlain by approximately 60% morainal plain and up to 40% glaciolacustrine veneer		
'MvLgVFGp'	indicates at least 60% of the area is underlain by morainal veneer, with up to 40% glaciolacustrine veneer and less than 15% glaciofluvial plain		
'LgP/M'	indicates more than 60% of the area is underlain by a glaciolacustrine plain, with less than 15% moraine		
Stratigraphic Sequence			
Where materials of different origins or textures are known to be superimposed or can be confidently inferred, the sequence is indicated in conventional order using vertical separators, such as:			
'sLgV Mp'	indicates sandy glaciolacustrine veneer deposited on morainal plain		
Transitional Association			
Locally, two or more terrain units are juxtaposed by reason of related origin, temporal sequence or ambiguous geomorphological distinction. In the last case, both components may or may not be present. Such situations are identified by a compound designation marked by a hyphen. Examples are:			
'FGz-Lgz'	indicates ice-contact delta indistinguishable from glaciolacustrine delta		
'Lg-LGL'	indicates glaciolacustrine indistinguishable from littoral and nearshore glaciolacustrine sediment		
Morphological Overprint			
Where a sequence of geomorphological processes has produced a multi-aspect or compound terrain fabric, the geomorphological modifier suffixes are appended in the inferred order of superposition. 'Mpy' indicates a morainal plain has been moulded into ridges and finally dissected by streams. 'FGpfr' indicates a glaciofluvial plain that includes discontinuous hummocks and ridges.			
Methodology			
The Alberta Geological Survey conducted surficial geology field mapping in the area during 2011. Observations made during field mapping were combined with the interpretation of Light Detection and Ranging (LIDAR) bare-earth data (Figure 1), digital orthorectified aeriophotographs (1:60 000 scale, taken during 1984), and image classification of peatlands from Landsat 7 multispectral data (Natural Resources Canada, 2004). The LIDAR digital elevation model was used to delineate landforms through shaded-relief images created from three illumination directions. The shaded relief shown as an underlay on the main map was produced by using shaded-relief (315° illumination azimuth, 45° declination) and slope-gradient images.			
Acknowledgements			
B. Hogberg, C. Heikinen, and K. McKay assisted with the fieldwork. K. McKay completed the digital cartography and GIS. Spatial Data Warehouse Ltd. provided the base data. D. J. Utting provided constructive comments that improved this map.			
References			
Andriashek, L.D. (2001): Surficial geology of the Wapiti area, Alberta (NTS 83L); Alberta Energy and Utilities Board, EUB/AGS Map 239, scale 1:250 000.			
Atkinson, N. and Pawley, S.M. (2013): Surficial geology of the Crooked Lake area (NTS 83K/SW); Energy Resources Conservation Board, ERCB/AGS Map 563, scale 1:100 000.			
Evans, D.J.A. (2005): Glacial landscapes; Oxford University Press, New York, New York, 532 p.			
Natural Resources Canada (2004): Landsat 7 orthorectified imagery over Canada; Natural Resources Canada, Earth Observation Imagery.			
Roed, M.A. (1970): Surficial geology of Edson (NTS 83F); Research Council of Alberta, Alberta Geological Survey, Map 139, scale 1:250 000.			
Utting, D.J. (2012): Surficial geology of the Meekwap Lake area (NTS 83K/NE); Energy Resources Conservation Board, ERCB/AGS Map 559, scale 1:100 000.			
Utting, D.J. (2013): Surficial geology of the Fox Creek area (NTS 83K/SE); Energy Resources Conservation Board, ERCB/AGS Map 562, scale 1:100 000.			




Alberta Geological Survey
7801-422-1107
www.ags.gov.ab.ca
Published 2013
ISBN 978-1-4001-0098-1

Map 564
Surficial Geology of the Little Smoky Area (NTS 83K/NW)
Geology by: S.M. Pawley and N. Atkinson



Projection: Universal Transverse Mercator
Datum: North American Datum, 1983



Recommended Reference Format
Pawley, S.M. and Atkinson, N. (2013): Surficial geology of the Little Smoky area (NTS 83K/NW); Energy Resources Conservation Board, ERCB/AGS Map 564, scale 1:100 000.