


This is a common map legend for the surficial geology of northern Alberta. Coloured legend blocks indicate map units that appear on this map. Not all map symbols shown in the legend necessarily appear on this map.


UNIT	UNIT NAME	DESCRIPTION AND GENESIS
QUATERNARY		
HLOCENE		
A	Anthropogenic materials	Artificially made ground or geological materials that have been disturbed by human activity, such that their physical properties (e.g. structure, cohesion, compaction) have been drastically altered.
O	Organic deposits	Undifferentiated peat (woody to fibrous muck) occurring in wetlands; commonly underlain by fine-grained, poorly drained glaciolacustrine deposits; includes marshes, swamps, bogs and fens.
OB	Bog peat	Occurs in a peatland with a fluctuating water table and commonly a raised surface; peatland surface is dominated by sphagnum mosses, heath shrubs and short, stunted trees.
OF	Fen peat	Occurs in peatland which receives water from slowly flowing streams and groundwater, with the water table lying at the land surface; peatland surface is dominated by sedges, with grasses and reeds near local pools, and is sparsely treed.
C	Colluvial deposits	Materials that have reached their present position as a result of direct, gravity-induced movement, commonly occurs as slope and slump deposits confined to valley slopes and floors; includes pre-existing bedrock, till, glaciolacustrine, glacioluvial and eolian sediments, generally poorly sorted.
F	Fluvial deposits	Sediments transported and deposited by streams and rivers; synonymous with alluvium. Includes well-sorted stratified sand, gravel, silt, clay and organic sediments occurring in channel and overbank deposits (e.g., postglacial floodplains, terraces, fans and deltas).
L	Lacustrine deposits	Sediments deposited in and adjacent to recent and modern lakes; includes offshore sand, silt and clay, minor organic deposits; may also include minor littoral (nearshore) beaches and bars composed of sand, silt and minor gravel.
E	Eolian deposits	Wind deposited sediments; comprise well-sorted, medium- to fine-grained sand and minor silt, generally massive to locally cross-bedded or ripple-laminated; includes both active and vegetated dunes and sand sheets.
PLEISTOCENE		
LG	Glaciolacustrine deposits	Primarily fine-grained, distal sediments deposited in or along the margins of glacial lakes, including sediments released by the melting of floating ice. Includes laminated (rhythmically bedded) to massive fine sand, silt and clay, and may contain ice-rafted debris.
LGL	Littoral and nearshore sediments	Massive to stratified, well-sorted silty sand, pebbly sand and minor gravel; occurs in beaches, bars, spits and deltaic foresets deposited during regression and lowering of glacial lakes.
FG	Glacioluvial deposits	Sediments deposited by glacial meltwater streams as subaerial or subaqueous outwash. Includes sand and gravel, often stratified, minor silt, and may show evidence of ice melting (slumped structures). Features include meltwater channels, kettle holes, terraces and minor ice-contact sediments.
FGI	Ice-contact sediments	Sediments deposited by meltwater streams flowing either in direct contact with the ice margin (kame terraces) or within and/or under glacial ice (eskers, crevasse ridges). Includes massive to stratified, poor to moderately sorted, coarse-grained sediments (predominantly pebble gravel and coarse-grained sand, locally till) and may show evidence of ice melting (slumped structures).
M	Moraine	Diamictic (till) deposited directly by glacial ice and consisting of a mixture of clay, silt, sand and minor pebbles, cobbles and boulders. Locally this unit may contain blocks of bedrock, pre-existing stratified sediment and till, or lenses of glaciolacustrine and/or glacioluvial sediment.
MS	Stagnant ice moraine	Material resulting from the collapse and slumping of englacial and supraglacial sediment in response to the melting of buried stagnant ice at the ice margin; sediment is mainly diamictic, but locally includes stratified sediments of glaciolacustrine or glacioluvial origin. Characterized by low to high-relief hummocky topography.
MT	Ice-thrust moraine	Terrain formed from the glaciotectonic displacement of materials as blocks or rafts in a more or less intact state. Materials may include syngenetic till, as well as masses of pre-existing sediments and/or bedrock. Characterized by high to moderate relief and features include hill-hole pairs and glaciotectonic moraines.
MF	Fluted moraine	Glacially streamlined terrain; varies from alternating furrows and ridges to nearly equidimensional smoothed hills; all landforms parallel the local ice flow direction; includes flutes, drumlins and drumlinoids.
FP	Preglacial fluvial deposits	Sediments transported and deposited by streams and rivers prior to glaciation. This includes sand and gravel deposited in paleovalleys (i.e., preglacial floodplains, terraces, fans and deltas).
PRE-QUATERNARY		
RT	Unconsolidated fluvial gravels	Predominantly well-sorted, quartzite and chert gravel and cobbles; Cordilleran source, Paleogene to Neogene.
R	Bedrock	

SYMBOL LEGEND		BASEMAP LEGEND	
Permafrost; relic and/or active		River	
Meltwater channel (minor)		Lake	
Meltwater channel (minor, paleoflow direction known)		UTM, Zone 11 Grid	
Crevasse filling		Contour, intervals 50 metres	
Ice thrust ridge			
Esker (paleoflow direction unknown)			
Drumlinoid or streamlined landform			
Minor moraine ridge			
Major moraine ridge			

UNIT NOTATION		
Example: sandy GLACIOLACUSTRINE plain		
Textural modifier	s LG p Geomorphic unit	
Textural Modifier		
Textural characteristics may be applied to the terrain classification as a prefix based on field observations or by inference from distinctive genesis and/or morphology. When two modifiers are given, the second letter is the dominant texture, with the first letter indicating the secondary texture; i.e., sc for sandy clay.		
p	pebble	
g	gravel	
s	sand	
l	silt	
c	clay	
a	sand-silt-clay	
GENETIC & GEOMORPHOLOGICAL MODIFIERS		
c	crevasse fill	ice-contact ridges formed by the slumping of sediment into crevasse on the ice surface or the squeezing of till into fractures at the ice base
d	doughnut rings and ridges	circular hummocks with a central depression, plateau mounds and brain-like pattern ridges, low to moderate relief
e	eroded	planar surface eroded by glacial meltwater, often capped by a boulder lag and/or thin deposit of sand and gravel
g	gullied	slopes dissected by modern ravines created by intermittent runoff
h	hummock	assemblage of approximately equidimensional hills and hollows; moderate to high relief (commonly greater than 2 m)
k	collapse	depression, includes kettle holes, pitted morphology, thermokarst depressions, karst sinkholes
m	meander	sinuous curves, loops and oxbows produced as meltwater and modern streams shift their channels over time
p	plain	deposit greater than 2 m thick, commonly masks geomorphic pattern of underlying deposits; flat to gently rolling topography (commonly less than 2 m relief)
r	ridged	one or more parallel or subparallel, convex, linear morphological elements with a length-to-width ratio greater than 2;
s	slumped	landslide blocks, slope failure debris
t	terrace	a bench of either erosional or depositional origin that flanks the sides of floodplains, valleys and lakes; includes fluvial and glacioluvial terraces, shoreline terraces and antiplation terraces
u	undulating	low-relief rolling terrain; swell and swale topography
v	vener	thin mantle of unconsolidated sediment that is too thin to mask the minor irregularities of the surface of the underlying material; it ranges in thickness from 10 cm to 1 metre and may be discontinuous
Complex		
Where two or more classes of terrain are interspersed in a mosaic or repeating pattern on a scale too small to warrant meaningful differentiation, the proportion of each component in the combination is given in a two- or three-position designation set off by slashes denoting arbitrary percentage limits. Examples are:		
'Mp/LGv'	indicates the area is underlain by approximately 60% morainal plain and up to 40% glaciolacustrine veneer	
'Mv/LGv/FGp'	indicates at least 60% of the area is underlain by morainal veneer, with up to 40% glaciolacustrine veneer and less than 15% glacioluvial plain	
'LGP/M'	indicates more than 60% of the area is underlain by a glaciolacustrine plain, with less than 15% moraine	
Stratigraphic Sequence		
Where materials of different origins or textures are known to be superimposed or can be confidently inferred, the sequence is indicated in conventional order using vertical separators, such as:		
'sLGv Mp'		
Transitional Association		
Locally, two or more terrain units are juxtaposed by reason of related origin, temporal sequence or ambiguous geomorphological distinction. In the last case, both components may or may not be present. Such situations are identified by a compound designation marked by a hyphen. Examples are:		
'LG-LGL'	indicates glaciolacustrine indistinguishable from littoral and nearshore glaciolacustrine sediment	
Morphological Overprint		
Where a sequence of geomorphological processes has produced a multi-aspect or compound terrain fabric, the geomorphological modifier suffixes are appended in the inferred order of superposition. 'Mpry' indicates a morainal plain has been moulded into ridges and finally dissected by streams. 'FGpr' indicates a glacioluvial plain that includes discontinuous hummocks and ridges.		
Methodology		
The Alberta Geological Survey conducted surficial geology field mapping in the area during 2014. Observations made during field mapping were combined with the interpretation of Light Detection and Ranging (LiDAR) bare-earth data (Figure 1) and image classification of peatlands from Landsat 5 and Landsat 8 multispectral data. The LiDAR digital elevation model was used to delineate landforms through shaded-relief images created from three illumination directions. The shaded relief shown as an underlay on the main map was produced by fusing shaded-relief (315° illumination azimuth, 45° declination) and slope-gradient images.		
Acknowledgements		
S. Pawley and L. Atkinson performed the fieldwork. K. McKay completed the digital cartography and GIS. Government of Alberta provided the base data. D. Utting provided comments that improved this map.		
References		
Atkinson, N., Utting, D.J. and Pawley, S.M. (2014). Glacial landforms of Alberta; Alberta Energy Regulator, AER/AGS Map 604, scale 1:1 000 000.		
Fenton, M.M., Waters, E.J., Pawley, S.M., Atkinson, N., Utting, D.J. and McKay, K. (2013). Surficial geology of Alberta; Alberta Energy Regulator, AER/AGS Map 601, scale 1:1 000 000.		
Paulin, R.C. and Plooffs, A. (2007). Surficial geology of the La Crete area (NTS 84G/SE); Alberta Energy and Utilities Board, EUB/AGS Map 412 and Geological Survey of Canada, Open File 5526, scale 1:1 000 000.		



Map 576
Surficial Geology of the Mikkwa River Area (NTS 84G/NE)
Geology by: S.M. Pawley



Projection: Universal Transverse Mercator
Datum: North American Datum, 1983

