

This is a common map legend for the surficial geology of Alberta. Coloured legend blocks indicate map units that appear on this map. Not all map symbols shown in the legend necessarily appear on this map.

UNIT	UNIT NAME	DESCRIPTION AND GENESIS
QUATERNARY		
HOLOCENE		
A	ANTHROPOGENIC MATERIALS	Artificially made ground or geological materials that have been disturbed by human activity, such that their physical properties (e.g. structure, cohesion, compaction) have been drastically altered.
O	ORGANIC DEPOSITS	Undifferentiated peat (woody to fibrous muck) occurring in wetlands; commonly underlain by fine-grained, poorly drained glaciolacustrine deposits; includes marshes, swamps, bogs and fens.
OB	Bog peat	Occurs in a peatland with a fluctuating water table and commonly a raised surface; peatland surface is controlled by sphagnum mosses, heath shrubs and short, stunted trees.
OF	Fen peat	Occurs in peatland which receives water from slowly flowing streams and groundwater; with the water table lying at the land surface; peatland surface is dominated by sedges, with grasses and reeds near local scows, and is sparsely treed.
C	COLLUVIAL DEPOSITS	Materials that have reached their present position as a result of direct, gravity-induced movement; commonly occurs as slope and slump deposits confined to valley slopes and floors; includes pre-existing bedrock, till, glaciolacustrine, glaciolacustrine and eolian sediments; generally poorly sorted.
F	FLUVIAL DEPOSITS	Sediments transported and deposited by streams and rivers; synonymous with alluvium. Includes well-sorted stratified sand, gravel, silt, clay and organic sediments occurring in channel and overbank deposits (e.g., postglacial floodplains, terraces, fans and deltas).
L	LACUSTRINE DEPOSITS	Sediments deposited in and adjacent to recent and modern lakes; includes offshore sand, silt and clay; minor organic deposits; may also include minor littoral (nearshore) terraces and bars composed of sand, silt and minor gravel.
E	EOLIAN DEPOSITS	Wind deposited sediments; comprise well-sorted, medium- to fine-grained sand and minor silt; generally massive to locally cross-bedded or ripple-laminated; includes both active and vegetated dunes and sand sheets.
PLEISTOCENE		
LG	GLACIOLACUSTRINE DEPOSITS	Primarily fine-grained, distal sediments deposited in or along the margins of glacial lakes, including sediments released by the melting of floating ice. Includes laminated (rhythmically bedded) to massive fine sand, silt and clay, and may contain ice-raftered debris.
LGL	Littoral and nearshore sediments	Massive to stratified, well-sorted silt, sand, pebbly sand and minor gravel; occurs in beaches, bars, spits and deltaic foresets deposited during regression and lowering of glacial lakes.
LGLI	Ice-contact sediments	Sediments deposited in ice-walled lake plains along the margins of stagnant glacial ice. Characterized by low- to high-relief hummocky topography, including flat-topped hills. Typically comprise glaciolacustrine sediment, including laminated to massive fine sand, silt and clay in the central part of these features, and littoral sediments around their margins. Locally contain distinct resulting from the collapse, melt-out or slumping of supraglacial debris from the surrounding ice walls.
FG	GLACIOFLUVIAL DEPOSITS	Sediments deposited by glacial meltwater streams as subaerial or subaqueous outwash. Includes sand and gravel, often stratified, minor silt, and may show evidence of ice melting (slumped structures). Features include meltwater channels, kettle holes, terraces and minor ice-contact sediments.
FGI	Ice-contact sediments	Sediments deposited by meltwater streams flowing either in direct contact with the ice margin (kame terraces) or within and/or under glacial ice (eskers, crevasse ridges). Includes massive to stratified, poor to moderately sorted, coarse-grained sediments (predominantly pebble gravel and coarse-grained sand, locally till) and may show evidence of ice melting (slumped structures).
M	MORAINES	Diamictic (till) deposited directly by glacial ice and consisting of a mixture of clay, silt, sand and minor pebbles, cobbles and boulders. Locally, this unit may contain blocks of bedrock, pre-existing stratified sediment and till, or lenses of glaciolacustrine and/or glaciolacustrine sediment.
MS	Stagnant ice moraine	Material resulting from the collapse and slumping of englacial and supraglacial sediment in response to the melting of buried stagnant ice at the ice margin; sediment is mainly diamictic, but locally includes stratified sediments of glaciolacustrine or glaciolacustrine origin. Characterized by low- to high-relief hummocky topography.
MT	Ice-thrust moraine	Terrain formed from the glaciectonic displacement of materials as blocks or rafts in a more or less intact state. Materials may include syngenetic till, as well as masses of pre-existing sediments and/or bedrock. Characterized by high to moderate relief and features include hill-hole pairs and glaciectonic moraines.
MF	Fluted moraine	Glacially streamlined terrain; varies from alternating furrows and ridges to nearly equidimensional smoothed hills; all landforms parallel the local ice flow direction; includes flutes, drumlins and drumlinoids.
FP	PREGLACIAL FLUVIAL DEPOSITS	Sediments transported and deposited by streams and rivers prior to glaciation. The include sand and gravel deposited in paleovalleys (i.e., preglacial floodplains, terraces, fans and deltas).
PRE-QUATERNARY		
RT	UNCONSOLIDATED FLUVIAL GRAVELS	Predominantly well-sorted, quartzite and chert gravel and cobbles; Cordilleran source, Paleogene to Neogene.
R	BEDROCK	

SYMBOL LEGEND	BASEMAP LEGEND
Eolian forms: dune ridges	Primary road, paved
Escarpment	River
Meltwater channel (minor)	Lake
Meltwater channel (major)	10 Transverse Mercator Grid
Crevasse filling	Contour, intervals 100 metres
Ice-thrust ridge	City
Esker (paleoflow direction unknown)	Town
Esker (paleoflow direction known)	
Drumlinoid or streamlined landform	
Drumlinoid (ice flow direction known)	
Minor moraine ridge	
Major moraine ridge	
Buried features	

UNIT NOTATION
Example: sandy GLACIOLACUSTRINE plain

Textural Modifier
Textural characteristics may be applied to the terrain classification as a prefix based on field observations or by inference from distinctive genesis and/or morphology. When two modifiers are given, the second letter is the dominant texture, with the first letter indicating the secondary texture; i.e., sc for sandy clay.

GENETIC & GEOMORPHOLOGICAL MODIFIERS

c crevasse fill ice-contact ridges formed by the slumping of sediment into crevasses on the ice surface or the sequencing of till into fractures at the ice base

f fan gently sloping fan-shaped mass of detrital debris

g gullied slopes dissected by modern ravines created by intermittent runoff

h hummock assemblage of approximately equidimensional hills and hollows; moderate to high relief (commonly greater than 2 m)

k collapse depression, includes kettle holes, pitted morphology, thermokarst depressions, karst sinkholes

m meander sinuous curves, loops and oxbows produced as meltwater and modern streams shift their channels over time

p plain flat to gently rolling topography; deposit commonly greater than 2 m thick; may mask geomorphic pattern of underlying deposits

r ridged one or more parallel or subparallel, convex, linear morphological elements with a length-to-width ratio greater than 2; low to high relief

s slumped landslide blocks; slope failure debris

t terrace a bench of either erosional or depositional origin that flanks the sides of floodplains, valleys and lakes; includes fluvial and glaciolacustrine terraces, esker terraces and antipathway terraces

u undulating low-relief rolling terrain; swell and swale topography

v veneer thin mantle of unconsolidated sediment that is too thin to mask the minor irregularities of the surface of the underlying material; it ranges in thickness from 10 cm to 1 metre and may be discontinuous

Complex
Where two or more classes of terrain are interspersed in a mosaic or repeating pattern on a scale too small to warrant meaningful differentiation, the proportion of each component in the combination is given in a two- or three-position designation set off by slashes denoting arbitrary percentage limits. Examples are:
"MpLGV" indicates the area is underlain by approximately 60% morainal plain and up to 40% glaciolacustrine veneer
"MvLGV/FG" indicates at least 50% of the area is underlain by morainal veneer, with up to 40% glaciolacustrine veneer and less than 15% glaciolacustrine plain
"LGV/IM" indicates more than 60% of the area is underlain by a glaciolacustrine plain, with less than 15% moraine

Stratigraphic Sequence
Where materials of different origins or textures are known to be superimposed or can be confidently inferred, the sequence is indicated in conventional order using vertical separators, such as:
"sLGV/MP" indicates sandy glaciolacustrine veneer deposited on morainal plain


Transitional Association
Locally, two or more terrain units are juxtaposed by reason of related origin, temporal sequence or ambiguous geomorphological distinction. In the last case, both components may or may not be present. Such situations are identified by a compound designation marked by a hyphen. Examples are:
"LG-LGL" indicates glaciolacustrine indistinguishable from littoral and nearshore glaciolacustrine sediment

Morphological Overprint
Where a sequence of geomorphological processes has produced a multi-aspect or compound terrain fabric, the geomorphological modifier suffixes are appended in the inferred order of superposition. "Mgp" indicates a morainal plain has been moulded into ridges and finally dissected by streams. "FGp" indicates a glaciolacustrine plain that includes discontinuous hummocks and ridges.

Acknowledgements
Government of Alberta provided the base data. N. Atkinson, M. Fenton, S. Pawley and D. Utting provided comments that improved this map.

References
Alberta Agriculture and Rural Development (2014). Agricultural regions of Alberta Soil Inventory Database (AGRID). Version 4.0. ESRRI ArcGIS geodatabase.
URL: <http://www.agric.gov.ab.ca/department/9693000.nsf/Alltag14652m2adaa0ARASID40> (August 2014).
Alberta Environment and Parks (2015). Provincial DEM, digital topographic data. AtlasUS.
URL: <http://www.atlas.com/products/terrain/DEM.html>, 25 m cell. Esri Grid format (February 2016).
Atkinson, N., Utting, D.J. and Pawley, S.M. (2014). Glacial landforms of Alberta, Canada. Alberta Energy Regulator, AER/AGS Map 604, scale 1:1 000 000.
Budney, H.D., Edwards, W.A.D., Berezniuk, T. and Bullock, L. (2004a). Sand and gravel deposits with aggregate potential, Drumheller, Alberta (NTS 82P). Alberta Energy and Utilities Board, EUB/AGS Map 272, scale 1:250 000.
Budney, H.D., Edwards, W.A.D., Berezniuk, T. and Bullock, L. (2004b). Sand and gravel deposits with aggregate potential, Gleichen, Alberta (NTS 82J). Alberta Energy and Utilities Board, EUB/AGS Map 274, scale 1:250 000.
Budney, H.D., Edwards, W.A.D., Berezniuk, T. and Bullock, L. (2004c). Sand and gravel deposits with aggregate potential, Kamaska Lakes, Alberta (NTS 82I). Alberta Energy and Utilities Board, EUB/AGS Map 268, scale 1:250 000.
Burke, J.B., Brennan, T.A. and Slogren, D.B. (2015). The role of sediment supply in esker formation and ice tunnel evolution. Quaternary Science Reviews, v. 115, p. 50-77.
Edwards, W.A.D., Budney, H.D., Berezniuk, T., Cadin, M.J. and Stoyko, G. (2004). Sand and gravel deposits with aggregate potential, Lethbridge, Alberta (NTS 82H). Alberta Energy and Utilities Board, EUB/AGS Map 287, scale 1:250 000.
Fenton, M.M., Waters, E.J., Pawley, S.M., Atkinson, N., Utting, D.J. and McKay, K. (2013). Surficial geology of Alberta. Alberta Energy Regulator, AER/AGS Map 601, scale 1:1 000 000.
Jackson, L.E., Jr. (1988a). Surficial geology, Graman, Alberta. Geological Survey of Canada, Map 1929A, scale 1:50 000.
Jackson, L.E., Jr. (1988b). Surficial geology, Langford Creek, Alberta. Geological Survey of Canada, Map 1927A, scale 1:50 000.
Jackson, L.E., Jr. (1988c). Surficial geology, Maycroft, Alberta. Geological Survey of Canada, Map 1928A, scale 1:50 000.
Jackson, L.E., Jr. (1988d). Surficial geology, Stinson Creek, Alberta. Geological Survey of Canada, Map 1926A, scale 1:50 000.
Jackson, L.E., Jr. (1988e). Surficial geology, Turner Valley, Alberta. Geological Survey of Canada, Map 1925A, scale 1:50 000.
Lebow, E.R. (1998a). Surficial geology, Brocket, Alberta. Geological Survey of Canada, Map 1931A, scale 1:50 000.
Lebow, E.R. (1998b). Surficial geology, Pincher Creek, Alberta. Geological Survey of Canada, Map 1933A, scale 1:50 000.
Little, E.C. (1998a). Surficial geology, Cardston, Alberta. Geological Survey of Canada, Map 1936A, scale 1:50 000.
Little, E.C. (1998b). Surficial geology, Raley, Alberta. Geological Survey of Canada, Map 1934A, scale 1:50 000.
Moran, S.R. (1988). Surficial geology of the Calgary urban area, Terrain Sciences Department, Alberta Research Council, Edmonton, AB.
Shetsen, I. (1981). Surficial geology, Lethbridge, Alberta (NTS 82H, 82I). Alberta Research Council, Alberta Geological Survey, Map 209, scale 1:250 000.
Shetsen, I. (1987). Quaternary geology, Southern Alberta, Terrain Sciences Department, Alberta Research Council, Map 207, scale 1:500 000.
Stalker, A.MacS. (1957). Surficial geology, High River, West of Fourth Meridian, Alberta. Geological Survey of Canada, Map 14-1957, scale 1:253 440.
Stalker, A.MacS. (1959). Surficial geology, Fort MacLeod, West of Fourth Meridian, Alberta. Geological Survey of Canada, Map 21-1959, scale 1:253 440.
Stalker, A.MacS. (1962). Surficial geology, Lethbridge (east half), West of Fourth Meridian, Alberta. Geological Survey of Canada, Map 41-1962, scale 1:253 440.
Stalker, A.MacS. (1963). Surficial geology, Blood Indian Reserve, West of Fourth Meridian, Alberta. Geological Survey of Canada, Map 40-1963, scale 1:93 390.
Stalker, A.MacS. (1965). Surficial geology, Bassano (Gleichen, east half), West of Fourth Meridian, Alberta. Geological Survey of Canada, Map 5-1965, scale 1:253 440.
Stalker, A.MacS. and Crisp, B.G. (1973). Surficial geology, Drumheller, West of Fourth Meridian, Alberta. Geological Survey of Canada, Map 1386A, scale 1:250 000.

Recommended Reference Format
Hartman, G.M.D. (2016). Surficial geology of the Calgary-Lethbridge corridor, Alberta (NTS 82O, 82P, 82J, 82I, 82G and 82H). Alberta Energy Regulator, AER/AGS Map 579, scale 1:250 000.



Alberta Geological Survey
(780) 638-4341
www.agr.ab.ca

Published 2016
ISBN 978-1-4601-0154-4

Map 579
Surficial Geology of the Calgary-Lethbridge Corridor
NTS 82O, 82P, 82J, 82I, 82G and 82H
Geology Compiled by: G.M.D. Hartman

